To discuss with your Health Care Provider

Florida Blue 🛂 🗓



Adult (age 19+) Wellness Schedule

Be sure to review your plan benefits to determine your costs for these services

| Routine Health Guide | | | | | | | |
|---|---|--|--|--|--|--|--|
| Annual Wellness and Routine Check-up | Annually: Discuss related screening with your doctor. | | | | | | |
| Obesity Screening: Diet/Physical Activity/BMI Counseling | Annually. | | | | | | |
| Vision and Dental Exam (These services may not be covered by your medical benefits plan. Check your plan documents.). | Annually: Discuss with your doctor | | | | | | |
| Recommended Diagnostic Checkups and Scree | enings for At-Risk Patients | | | | | | |
| Abdominal Aortic Aneurysm (AAA) Check | One-time screening for ages 65 to 75 who have ever smoked or have a family history of AAA | | | | | | |
| Bone Mineral Density Screening and prescribed medication for Osteoporosis | Women beginning at age 65 or older; and in younger women who have an increased risk | | | | | | |
| Cholesterol Screening | Age 35+; Age 20-35 at risk Annually: All Men; Age 45+; Age 45 at risk Annually: All Women | | | | | | |
| Colorectal Cancer Screening and Counseling | Age 45-75; Colonoscopy or fecal occult blood test or sigmoidoscopy | | | | | | |
| Mammogram | Women should have a baseline mammogram age 35-40. Thereafter, every two years age 40-50; every year age 50+. At any age if recommended by physician based on risk factors. | | | | | | |
| Pap Test/Pelvic Exam | Women age 21-29 should have a Pap Test every 3 years. Women age 30-65 should have a Pap Test alone every 3 years or combined with HPV testing every 5 years. Women ages 65+ should discuss with their doctor. | | | | | | |
| HIV and other Sexually Transmitted Infections (STIs) Screening and Counseling | As indicated by history and/or symptoms. Discuss with your doctor behavioral risks. | | | | | | |
| Lung Cancer Screening and Counseling | Ages 50-80; 20 pack smoker history, current smoker/quit within past 15 years | | | | | | |
| Prostate Cancer Screening | Discuss with your doctor | | | | | | |
| Skin Cancer Screening | Discuss with your doctor | | | | | | |

Live a Healthy Lifestyle

- Get your annual wellness exam to review your overall health and keep follow-up visits with your doctor.
- Find out if you are at risk for health conditions such as diabetes, high cholesterol and high blood pressure.
- Get your vaccines, preventive screenings and labs.
- Human Papillomavirus (HPV) vaccine 3 dose series is recommended for men and women ages 19 through 26 years if not previously vaccinated prior to age 13.
- Talk with your doctor about the medications and over-thecounter/vitamins you are taking to reduce side effects and interactions.
- Get a Flu Vaccine every year to prevent illness and related hospitalizations.
- Get a COVID-19 vaccine to prevent severe illness and related hospitalizations. Immunocompromised people should consult their physician on the need for an additional mRNA vaccine dose.


Sources: These guidelines are recommendations from the following organizations and were not developed by Florida Blue. www.ahrq.gov, www.cdc.gov

To discuss with your Health Care Provider

Adult (age 19+) Wellness Schedule
Be sure to review your plan benefits to determine your costs for these services

| Guidance | |
|---|---|
| Screen/Counseling: Depression, Obesity, Tobacco, Alcohol, Substance Abuse and Pregnancy | Every visit, or as indicated by your doctor |
| Fall Risk/Unintentional Injury/Domestic Violence Prevention/Seat Belt Use | Discuss exercise and home safety with your doctor |
| Medication List (including over-the-counter and vitamins) for potential interactions | Every visit, or as indicated by your doctor |
| Advance Directives/Living Will | Annually |

| Immunizations* (Routine Recommendations | |
|---|---|
| Tetanus, Diphtheria, Pertussis (Td/Tdap) | Ages 19+: Tdap vaccine once, then a Td booster every 10 years |
| Flu (Influenza) | Annually during flu season |
| Pneumococcal PCV13 and PPSV23 | Ages 19-64: if risk factors are present; Ages 65+: 1-2 doses (per CDC); Ages 50+:1 dose (Florida Blue Benefits) |
| Shingles (Zoster) | Ages 50+: 2 doses Shingrix |
| Haemophilus Influenzae Type b (HIB) Hepatitis A, Hepatitis B, Meningococcal | Ages 19+: if risk factors are present |
| Human Papillomavirus (HPV), Measles/Mumps/Rubella (MMR), Varicella (Chickenpox) & Hepatitis C (HCV) Infection Screening | Physician recommendation based on past immunization or medical history |
| COVID-19 | Recommended for adults ages 19 and older within the scope of the authorization/approval for the particular vaccine. |





Your local Blue Cross Blue Shield

We're here to help: Call

Customer Service 1-800-FLA-BLUE (1-800-352-2583) TTY/TDD Call 711

Care Consultant Team 1-888-476-2227

Go to floridablue.com

Visit a Florida Blue Center

Go to floridabluecenters.com for locations or call 1-877-352-5830

Sources: These guidelines are recommendations from the following organizations and were not developed by Florida Blue. www.ahrq.gov, www.cdc.gov

To discuss with your Health Care Provider

Florida Blue 🛂 🗓

Your local Blue Cross Blue Shield

Children & Adolescents (Birth – 18 years of age) Wellness Schedule

| Routine Health Guide | | | | | | |
|--|--|--|--|--|--|--|
| Wellness Exam and Autism/Development Behavioral Assessment | Newborn up to age 3: Frequent Wellness Check- ups; Age 3-18: Annual Wellness Check-up | | | | | |
| Body Mass Index (BMI): Height and Weight | Every visit, BMI beginning at age 2 | | | | | |
| Blood Pressure | Annually, beginning at age 3 | | | | | |
| Hearing/Dental/Vision Screenings (These services may not be covered by your medical benefits plan. Check your plan documents.) | Hearing: Newborn then annually beginning at Age 4; Dental: Regularly, beginning at age 1; Vision: Annually, beginning at age 3 | | | | | |
| Recommended Screenings for At-Risk Patients | | | | | | |
| Cholesterol Screening | Annually, beginning at age 2 | | | | | |
| Lead test, TB, Sickle Cell and Blood Sugar | As indicated by history and/or symptoms | | | | | |
| HIV and other Sexually Transmitted Infections (STIs) Screening and Counseling | Discuss with your doctor based on behavioral risks | | | | | |
| Skin Cancer Screening | Discuss with your doctor | | | | | |
| Guidance | | | | | | |
| Injury/Violence Prevention | Annually, more often if indicated by your doctor | | | | | |
| Diet/Physical Activity/Emotional Well-Being Counseling | Every visit | | | | | |
| Tobacco/Alcohol/Substance Abuse/Depression/Pregnancy Screening and Counseling | Every visit starting at age 11, earlier if indicated by your doctor | | | | | |

Are your children up-todate with vaccinations?

Getting the recommended sequence of vaccinations is always a good idea to protect your child from illnesses from birth to 18 years of age. Most of these vaccinations require additional doses or boosters over time. As children grow up to become teenagers, they may come in contact with different diseases. Here are vaccines that can help protect your preteen or teen from these other illnesses and infections:

Tdap Vaccine

Age 11 or 12. Protects against tetanus (lock jaw), diphtheria and acellular pertussis (whooping cough). This is a booster shot of the same vaccine given during early childhood.

Meningococcal Vaccine

(MCV4) Two doses beginning at 11 or 12 years, with a booster dose at age 16. (MenB) Two doses beginning at ages 16-18. Protects against meningitis, sepsis (a blood infection) and other meningococcal diseases. Children with higher risk factors may need additional doses.

Contiunued next page

Sources: These guidelines are recommendations from the following organizations and were not developed by Florida Blue. www.ahrq.gov, www.cdc.gov

To discuss with your Health Care Provider

| Immunizations* | Birth | 1 month | 2 months | 4 months | 6 months | 12 months | 15 months | 18 months | 24 months | 4 - 6 years | 11 - 12 years | 13 - 14 years | 15 years | 16 - 18 years | | |
|--|----------|------------|-------------|-------------|--|--------------|--------------|--------------|--------------|----------------|------------------|------------------|-------------|------------------|--|--|
| Hepatitis A | | | | | 2 dose series, 12-23 mo | | | | onths | | | | | | | |
| Hepatitis B | 1st dose | 2nd | dose | | 3rd dose | | | | | | | | | | | |
| Diphtheria, Tetanus, Pertussis (DTaP) | | | 1st dose | 2nd dose | 3rd dose | 4th dose | | 4th dose | | ose | | 5th dose | | | | |
| Tetanus, Diphtheria, Pertussis (Tdap) | | | | | | | | | | | 1st dose | | | | | |
| Haemophilus Influ- enzae Type b (Hib) | | | 1st dose | 2nd dose | 3rd or 4th dose** | | | | | | | | | | | |
| Inactivated Poliovirus | | | 1st dose | 2nd dose | 3rd dose | | | | 4th dose | | | | | | | |
| Measles, Mumps, Rubella (MMR)*** | | | | | 1st dose | | | | 2nd dose | | | | | | | |
| Varicela | | | | | 1st dose | | | | 2nd dose | | | | | | | |
| Pneumococcal PCV13 and PPSV23 | | | 1st dose | 2nd dose | 3rd dose 4th dose | | | | | | | | | | | |
| Flu (Influenza) | | | | | 6 months through 8 years 1 or 2 doses; 9 years and older 1 dose only | | | | | | | | | | | |
| Rotavirus | | | 1st dose | 2nd dose | 3rd dose** | | | | | | | | | | | |
| Meningococcal | | | | | | | | | | | 1st dose | | | booster | | |
| Human Papillomavirus (HPV) | | | | | | | | | | | 3 dose series | | | | | |
| Covid 19 | | | | | 12 and older per CDC guidelines f specific vaccine | | | | | | | lines for | | | | |

^{*}These are routine immunizations based upon cdc.gov recommendations. Range of recommended ages for catch-up or certain high-risk groups is at the doctor's discretion based on the member's family history and personal risk factors.



Your local Blue Cross Blue Shield

Human Papillomavirus

(HPV) Vaccine two or three doses over six months, beginning at ages 11–12. (Two doses if started before 15th birthday or three if started on or after 15). Protects boys and girls against HPV, which can lead to cancers and genital warts.

Flu Vaccine

Every year after six months of age. Protects individuals from getting the influenza virus.

COVID-19 Vaccine

The American Academy of Pediatrics (AAP) recommends the COVID-19 vaccination for all children and adolescents 12 years of age and older who do not have contraindications using a COVID-19 vaccine authorized for use for their age. Any authorized COVID-19 vaccine appropriate by age and health status can be used for COVID-19 vaccination in children and adolescents.

Keep your teens safe from preventable, painful and life-threatening diseases by staying in touch with your pediatrician's office or health clinic. Be sure to verify your benefits for preventive services.

Sources: These guidelines are recommendations from the following organizations and were not developed by Florida Blue. www.ahrq.gov, www.cdc.gov

Florida Blue is an Independent Licensee of the Blue Cross and Blue Shield Association.

^{**} Dosages determined by doctor with the type of brand vaccine used.

^{***} Those in an outbreak area, aged ≥12 months who previously received ≤ 2 doses should receive a dose of mumps-virus containing vaccine.